

THE FEICHTINGER CONJECTURE FOR WAVELET FRAMES, GABOR FRAMES AND FRAMES OF TRANSLATES

MARCIN BOWNIK AND DARRIN SPEEGLE

ABSTRACT. The Feichtinger conjecture is considered for three special families of frames. It is shown that if a wavelet frame satisfies a certain weak regularity condition, then it can be written as the finite union of Riesz basic sequences each of which is a wavelet system. Moreover, the above is not true for general wavelet frames. It is also shown that a sup-adjoint Gabor frame can be written as the finite union of Riesz basic sequences. Finally, we show how existing techniques can be applied to determine whether frames of translates can be written as the finite union of Riesz basic sequences. We end by giving an example of a frame of translates such that any Riesz basic subsequence must consist of highly irregular translates.

1. INTRODUCTION AND STATEMENTS OF MAIN RESULTS

A frame is a collection of elements $\{e_i : i \in I\}$ in a Hilbert space \mathcal{H} such that there exist positive constants A and B such that for every $h \in \mathcal{H}$,

$$A\|h\|^2 \leq \sum_{i \in I} |\langle h, e_i \rangle|^2 \leq B\|h\|^2.$$

A frame $\{e_i : i \in I\}$ is bounded if

$$\inf_{i \in I} \|e_i\| > 0.$$

(Note that it is automatic that $\sup_{i \in I} \|e_i\| < \infty$.) A sequence $\{e_i : i \in I\}$ is said to be a Riesz basic sequence if it is a Riesz basis for its closed linear span, i.e., there exist $K_1, K_2 > 0$ such that for every family of scalars $\{a_i : i \in I\}$

$$K_1 \sum_{i \in I} |a_i|^2 \leq \left\| \sum_{i \in I} a_i e_i \right\|^2 \leq K_2 \sum_{i \in I} |a_i|^2.$$

Note that for Riesz basic sequences we always have

$$0 < \inf_{i \in I} \|e_i\| \leq \sup_{i \in I} \|e_i\| < \infty.$$

With this notation, one can state the Feichtinger conjecture:

Conjecture 1.1. (*Feichtinger*) *Every bounded frame can be written as the finite union of Riesz basic sequences.*

Date: May 12, 2006.

2000 Mathematics Subject Classification. Primary 42B25, 42B35, 42C40.

Key words and phrases. frame, Riesz basic sequence, wavelets, Gabor system, frames of translates.

The first author was partially supported by the NSF grant DMS-0200080.

The Feichtinger conjecture has received a fair amount of recent interest [6, 7, 13], and it has been shown to be related to several famous open problems in analysis. In particular, in [6], it is shown that the conjectured generalization of the Bourgain-Tzafriri restricted invertibility theorem is equivalent to the Feichtinger conjecture. It is also shown that the paving conjecture of Bourgain-Tzafriri (which is known to be equivalent to the Kadison-Singer conjecture) implies the Feichtinger conjecture. For precise statements of these conjectures and references, see [6].

In this paper, we consider the Feichtinger conjecture for special systems of frames such as wavelet and Gabor frames and frames of translates. In particular, we are investigating a stronger version of the Feichtinger conjecture, which asks whether a bounded frame of a certain type (such as wavelet, Gabor, or frame of translates) can be decomposed as the finite union of Riesz sequences of the same type.

The answer to this question depends on the type of frame at hand. Namely, we prove that wavelet frames satisfying certain weak decay and regularity conditions satisfy the stronger Feichtinger property, whereas wavelet frames without this additional regularity condition in general do not satisfy this property. In contrast, we show that a significant class of Gabor systems satisfying a certain compatibility condition on their generating lattices always have this property regardless of any regularity conditions. Finally, for frames of translates the situation with regard to the stronger Feichtinger property is the same as for wavelets. Nevertheless, we show that the Feichtinger conjecture leads to some interesting number theoretic problems involving the quantitative versions of van der Waerden's theorem.

The plan of this paper is as follows. In Section 2, we introduce the notion of localization for wavelet frames associated with general expansive dilations and we show that for such frames the stronger Feichtinger conjecture holds. We prove that this result is optimal in the sense that it does not hold for general wavelet frames without a localization property. In Section 3 we show that every sup-adjoint Gabor frame can be written as the finite union of Riesz basic sequences, generalizing results of Ron and Shen [21] and Casazza et al. [6].

In Section 4, we study frames of subspaces of $L^2(\mathbb{R})$ of the form $\mathcal{S}_f = \{f(x+k) : k \in \Gamma\}$. We show that if $f(x)$ has good decay and \mathcal{S}_f is a frame of a subspace, then \mathcal{S}_f can be written as the finite union of Riesz basic sequences. Then, we consider the following

Question 1.2. *Suppose $\hat{f} = \chi_U$, where $U \subset \mathbb{R}$, and $\mathcal{S}_f = \{f(x+k) : k \in \mathbb{Z}\}$ is a frame for its closed linear span. Can \mathcal{S}_f be written as a finite union of Riesz basic sequences?*

Most interesting in this section is an example of a function $f(x)$ of the type $\hat{f} = \chi_U$ such that \mathcal{S}_f is a frame of a subspace, but such that if $N \subset \mathbb{Z}$ has arbitrarily long arithmetic progressions of a certain type, then $\{f(x+k) : k \in N\}$ is not a Riesz basic sequence. In particular, a random subset of \mathcal{S}_f almost surely is not a Riesz basic sequence, see Theorem 4.13 for details. This example is also compared to quantitative versions of van der Waerden's Theorem to conclude the section. Note that there is a close connection between the Feichtinger conjecture in this setting and

the *paving problem* for Laurent operators. This connection is also briefly explained in Section 4.

1.1. Some background tools. We start by recalling basic definitions and properties of non-isotropic Euclidean spaces associated with general expansive dilations. A real $n \times n$ matrix A is an *expansive matrix*, often called a *dilation*, if $\min_{\lambda \in \sigma(A)} |\lambda| > 1$, where $\sigma(A)$ is the set of all eigenvalues (the spectrum) of A . A fundamental notion in our study is a quasi-norm ρ_A associated with A .

Definition 1.1. A quasi-norm associated with an expansive matrix A is a measurable mapping $\rho_A : \mathbb{R}^n \rightarrow [0, \infty)$ satisfying

$$(1.1) \quad \begin{aligned} \rho_A(x) &> 0, && \text{for } x \neq 0, \\ \rho_A(Ax) &= |\det A| \rho_A(x) && \text{for } x \in \mathbb{R}^n, \\ \rho_A(x+y) &\leq H(\rho_A(x) + \rho_A(y)) && \text{for } x, y \in \mathbb{R}^n, \end{aligned}$$

where $H \geq 1$ is a constant.

Here, we will only list a few basic properties of quasi-norms ρ_A , which will be used subsequently. For more details we refer to [3, 18]. We recall that all quasi-norms associated to a fixed dilation A are equivalent, see [3, Lemma 2.4]. Moreover, there always exist a quasi-norm ρ_A , which is C^∞ on \mathbb{R}^n except the origin, see [18].

Proposition 1.3. *For any expansive matrix A and $\epsilon > 0$,*

$$\int_{B(0,1)} \rho_A(x)^{\epsilon-1} dx < \infty \quad \text{and} \quad \int_{\mathbb{R}^n \setminus B(0,1)} \rho_A(x)^{-1-\epsilon} dx < \infty.$$

Proposition 1.3 follows from

$$(1.2) \quad (1/c)r \leq |\{x \in \mathbb{R}^n : \rho_A(x) < r\}| \leq cr \quad \text{for all } r > 0.$$

Lemma 1.4. *Suppose A is expansive matrix, and λ_- and λ_+ are any positive real numbers such that $1 < \lambda_- < \min_{\lambda \in \sigma(A)} |\lambda|$ and $\lambda_+ > \max_{\lambda \in \sigma(A)} |\lambda|$. Let*

$$\zeta_- := \frac{\ln \lambda_-}{\ln |\det A|}, \quad \zeta_+ := \frac{\ln \lambda_+}{\ln |\det A|}.$$

Then for any quasi-norm ρ_A there exists a constant C such that,

$$(1.3) \quad C^{-1} \rho_A(x)^{\zeta_-} \leq |x| \leq C \rho_A(x)^{\zeta_+} \quad \text{if } \rho_A(x) \geq 1$$

and

$$(1.4) \quad C^{-1} \rho_A(x)^{\zeta_+} \leq |x| \leq C \rho_A(x)^{\zeta_-} \quad \text{if } \rho_A(x) \leq 1.$$

Furthermore, if A is diagonalizable over \mathbb{C} , then we may take $\lambda_- = \min_{\lambda \in \sigma(A)} |\lambda|$ and $\lambda_+ = \max_{\lambda \in \sigma(A)} |\lambda|$.

We will also need the following easy estimates

$$(1.5) \quad (1/c)\lambda_-^j |x| \leq |A^j x| \leq c\lambda_+^j |x| \quad \text{for } j \geq 0,$$

$$(1.6) \quad (1/c)\lambda_+^j |x| \leq |A^j x| \leq c\lambda_-^j |x| \quad \text{for } j \leq 0,$$

for some constant $c > 0$, where λ_- and λ_+ are the same as in Lemma 1.4.

For any $j \in \mathbb{Z}$ and $k \in \mathbb{Z}^n$, let $Q_{j,k} = A^{-j}([0, 1]^n + k)$ be the dilated cube, and $x_{Q_{j,k}} = A^{-j}k$ be its “lower-left corner”. Let

$$\mathcal{Q} = \{Q_{j,k} : j \in \mathbb{Z}, k \in \mathbb{Z}^n\}$$

be the collection of all dilated cubes. Given a function ψ on \mathbb{R}^n we define

$$(1.7) \quad \psi_Q(x) = |\det A|^{j/2} \psi(A^j x - k) = |Q|^{-1/2} \psi(A^j(x - x_Q)) \quad \text{for } Q = Q_{j,k} \in \mathcal{Q}.$$

Equivalently, if $Q = A^{-j}([0, 1]^n + k) \in \mathcal{Q}$ then $\psi_Q = D^j T_k \psi$, where D is a dilation operator $Df(x) = |\det A|^{1/2} f(Ax)$ and T_k is a translation operator $T_k f(x) = f(x - k)$.

2. AFFINE LOCALIZED FRAMES

Gröchenig [13] introduced a notion of localized frames and showed that the Feichtinger conjecture holds for this class of frames. While his definition works very well for systems generated by families of unitary operators with compatible actions on $L^2(\mathbb{R}^n)$, such as translations and modulations in the case of Gabor systems, there is a natural need to have a better adapted notion of localization for systems which are generated by families of operators with less compatible actions. The most prominent example of such system is an affine system generated by translates and dilates of a frame wavelet.

While, it seems that there is no single universal notion of frame localization working for all imaginable frames in $L^2(\mathbb{R}^n)$, we propose that the following definition gives a satisfactory answer in the case of wavelet systems. Our definition is motivated by the almost diagonal condition of Frazier and Jawerth [9, 10] from the study of isotropic Besov and Triebel-Lizorkin spaces and their anisotropic analogues for expansive dilations introduced by the first author and Ho [4].

Definition 2.1. We say that a collection $\{\Psi_Q : Q \in \mathcal{Q}\}$ is *affine localized* if there exists $C, \theta > 0$ such that,

$$(2.1) \quad |\langle \Psi_Q, \Psi_P \rangle| \leq C \kappa_{QP}(\theta) \quad \text{for all } Q, P \in \mathcal{Q},$$

where

$$\kappa_{QP}(\theta) = \left(1 + \frac{\rho_A(x_Q - x_P)}{\max(|P|, |Q|)}\right)^{-1-\theta} \min \left[\left(\frac{|Q|}{|P|}\right)^{1/2+\theta}, \left(\frac{|P|}{|Q|}\right)^{1/2+\theta} \right].$$

Remark 2.1. We remark that Ψ_Q should be only understood as a function indexed by $Q \in \mathcal{Q}$, which is not necessarily given by (1.7).

Definition 2.2. We say that $\Psi_Q(x)$ is a *molecule* supported near $Q \in \mathcal{Q}$ with $|Q| = |\det A|^{-j}$ and $j \in \mathbb{Z}$, if there exist $M > 1$ and $\delta > 0$ such that

$$(2.2) \quad |\Psi_Q(x) - \Psi_Q(y)| \leq \frac{|\det A|^{j(1/2+\delta)} \rho_A(x - y)^\delta}{(1 + \rho_A(A^j(x - x_Q)))^M}$$

for all $x, y \in \mathbb{R}^n$, $\rho_A(x - y) \leq |\det A|^{-j}$,

$$(2.3) \quad |\Psi_Q(x)| \leq \frac{|\det A|^{j/2}}{(1 + \rho_A(A^j(x - x_Q)))^M} \quad \text{for all } x \in \mathbb{R}^n,$$

$$(2.4) \quad \int_{\mathbb{R}^n} \Psi_Q(x) dx = 0.$$

We say that a collection $\{\Psi_Q\}_{Q \in \mathcal{Q}}$ is a *family of molecules*, if each Ψ_Q is a molecule supported near Q with the same $M > 1$ and $\delta > 0$.

Therefore, a molecule Ψ_Q must satisfy certain smoothness (2.2) and decay (2.3) conditions appropriately localized to a dilated cube Q in addition to the usual vanishing moment condition (2.4). We also mention that the smoothness condition (2.2) in Definition 2.2 can be replaced by

$$(2.5) \quad |\Psi_Q(x) - \Psi_Q(y)| \leq |\det A|^{j(1/2+\delta)} \rho_A(x-y)^\delta \sup_{\rho_A(z) \leq \rho_A(x-y)} (1 + \rho_A(A^j(x - z - x_Q)))^{-M}$$

to obtain an equivalent definition of a molecule supported near Q modulo a multiplicative constant independent of Q . A condition of the type (2.5) is more commonly used and it was introduced by Frazier and Jawerth for the dyadic dilation $A = 2Id$ in [10, Section 3]. To see the equivalence of (2.2) and (2.5) it suffices to make two observations. First, we have

$$\begin{aligned} 1 + \rho_A(A^j(x - x_Q)) &\leq \sup_{\rho_A(z) \leq \rho_A(x-y)} 1 + \rho_A(A^j(x - z - x_Q)) \\ &\leq (H + 1)(1 + \rho_A(A^j(x - x_Q))) \quad \text{for } \rho_A(x - y) \leq |\det A|^{-j}. \end{aligned}$$

Second, if $\rho_A(x - y) > |\det A|^{-j}$, then (2.3) immediately implies (2.5) with the right-hand side having an additional factor of 2.

Frequently, we will be interested in families of molecules generated by translates and dilates of a single function ψ . In this case, the following remark is relevant.

Remark 2.2. Suppose that a function $\psi(x)$ satisfies

$$(2.6) \quad |\psi(x) - \psi(y)| \leq \frac{\rho_A(x - y)^\delta}{(1 + \rho_A(x))^M} \quad \text{for } \rho_A(x - y) \leq 1,$$

$$(2.7) \quad |\psi(x)| \leq (1 + \rho_A(x))^{-M},$$

$$(2.8) \quad \int_{\mathbb{R}^n} \psi(x) dx = 0.$$

for some $\delta > 0$ and $M > 1$. Then, the affine system $\{\psi_Q : Q \in \mathcal{Q}\}$ given by (1.7) forms a family of molecules. Easy verification of this fact is left to the reader.

Also, for those who do not like the use of the quasi-norm ρ_A , one can write a slightly stronger version of conditions (2.6) and (2.7), where ρ_A is replaced by the standard Euclidean distance $|\cdot|$. Indeed, by Lemma 1.4 it suffices to assume that

$\psi(x)$ satisfies (2.8) and

$$(2.9) \quad |\psi(x) - \psi(y)| \leq \frac{|x - y|^\delta}{(1 + |x|)^{M/\zeta_-}} \quad \text{for } |x - y| \leq 1,$$

$$(2.10) \quad |\psi(x)| \leq (1 + |x|)^{-M/\zeta_-},$$

for some $\delta > 0$ and $M > 1$, to conclude that $\{\psi_Q : Q \in \mathcal{Q}\}$ given by (1.7) is a family of molecules in the sense of Definition 2.2.

We are now ready to state the key result showing that families of molecules must satisfy almost diagonal estimates.

Lemma 2.1. *Suppose $\{\Phi_Q\}_Q$ and $\{\Psi_Q\}_Q$ are families of molecules. Then there exist $C > 0$ and $\theta > 0$, such that*

$$|\langle \Psi_P, \Phi_Q \rangle| \leq C \kappa_{QP}(\theta) \quad \text{for all } Q, P \in \mathcal{Q}.$$

In particular, any family of molecules $\{\Psi_Q\}_Q$ is affine localized.

Lemma 2.1 was originally proved in the usual dyadic case by Frazier and Jawerth [10], in a much greater generality than needed here, for molecules with possible higher degrees of smoothness. In the case of general expansive dilations, the same was shown in [4]. Since the notion of smooth molecules in [4] is more restrictive than our Definition 2.2, we will give a proof of Lemma 2.1. We will need the following two elementary facts.

Lemma 2.2. *Suppose that $j \in \mathbb{Z}$, $j \geq 0$, $\theta > 0$, and $x \in \mathbb{R}^n$. Then there exists a constant $C > 0$, which is independent of j and x , such that*

$$(2.11) \quad \sum_{k \in \mathbb{Z}^n} (1 + \rho_A(A^{-j}k + x))^{-1-\theta} \leq C |\det A|^j.$$

Proof. First, note that Proposition 1.3 and a change of variables yield

$$\int_{\mathbb{R}^n} (1 + \rho_A(A^{-j}z + x))^{-1-\theta} dz \leq C |\det A|^j.$$

Next, it suffices to split the domain of integration into cubes $\mathbb{R}^n = \bigcup_{k \in \mathbb{Z}^n} (k + [0, 1]^n)$ and use the fact that for $z \in k + [0, 1]^n$,

$$\rho_A(A^{-j}z + x) \leq H(\rho_A(A^{-j}k + x) + \sup_{y \in [0, 1]^n} \rho_A(A^{-j}y)) \leq \tilde{C}(1 + \rho_A(A^{-j}k + x)).$$

□

Lemma 2.3. *Suppose that for some $\theta > 0$, $R > 1 + \theta$, and $i, j \in \mathbb{Z}$, $i \geq j$, and $x_0 \in \mathbb{R}^n$, we have*

$$(2.12) \quad |g(x)| \leq |\det A|^{j/2} (1 + \rho_A(A^j x))^{-R},$$

$$(2.13) \quad |g(x) - g(y)| \leq \frac{|\det A|^{j(1/2+\theta)} \rho_A(x-y)^\theta}{(1 + \rho_A(A^j x))^R} \quad \text{for } \rho_A(x-y) \leq |\det A|^{-j},$$

$$(2.14) \quad |h(x)| \leq |\det A|^{i/2} (1 + \rho_A(A^i(x-x_0)))^{-R},$$

$$(2.15) \quad \int_{\mathbb{R}^n} h(x) dx = 0.$$

Then there exists a constant $C > 0$ depending only on θ and R , such that

$$(2.16) \quad |(g * h)(x)| \leq C |\det A|^{-(i-j)(1/2+\theta)} (1 + \rho_A(A^j(x-x_0)))^{-R}.$$

Proof. Using translation and dilation, we may assume that $j = 0$ and $x_0 = 0$. Indeed, if g and h satisfy (2.12)–(2.14) for some $i_0 \geq j_0 \in \mathbb{Z}$ and $\tilde{x}_0 \in \mathbb{R}^n$, then $D^{-j_0}g(x)$ and $D^{-j_0}T_{-\tilde{x}_0}h(x)$ satisfy the corresponding conditions for $i = i_0 - j_0$, $j = 0$, and $x_0 = 0$. Thus, assuming Lemma 2.3 holds for $j = 0$ and x_0 ,

$$|(D^{-j_0}g * D^{-j_0}T_{-\tilde{x}_0}h)(x)| \leq C |\det A|^{-(i_0-j_0)(1/2+\theta)} (1 + \rho_A(x))^{-R}.$$

This will show the general case of Lemma 2.3, since

$$(D^{-j_0}g * D^{-j_0}T_{-\tilde{x}_0}h)(x) = (g * T_{-\tilde{x}_0}h)(A^{-j_0}x) = (g * h)(A^{-j_0}x + \tilde{x}_0).$$

Thus, assume that $j = 0$ and $x_0 = 0$, take any $x \in \mathbb{R}^n$ and decompose \mathbb{R}^n into 3 domains

$$\begin{aligned} D_1 &= \{y \in \mathbb{R}^n : \rho_A(y-x) < 1\}, \\ D_2 &= \{y \in \mathbb{R}^n : \rho_A(y-x) \geq 1 \quad \text{and} \quad \rho_A(y) \leq \rho_A(x)/(2H)\}, \\ D_3 &= \{y \in \mathbb{R}^n : \rho_A(y-x) \geq 1 \quad \text{and} \quad \rho_A(y) > \rho_A(x)/(2H)\}, \end{aligned}$$

where H is the constant of the quasi-subadditivity inequality.

By the vanishing moment condition (2.15) we have

$$|(g * h)(x)| \leq \int_{\mathbb{R}^n} |g(y) - g(x)| |h(x-y)| dy \equiv \int_{D_1} + \int_{D_2} + \int_{D_3}.$$

By (2.13), (2.14), the change of variables, and Proposition 1.3

$$\begin{aligned} \int_{D_1} &\leq |\det A|^{i/2} (1 + \rho_A(x))^{-R} \int_{D_1} \rho_A(x-y)^\theta (1 + \rho_A(A^i(x-y)))^{-R} dy \\ &\leq |\det A|^{i/2} (1 + \rho_A(x))^{-R} \int_{\mathbb{R}^n} \rho_A(y)^\theta (1 + \rho_A(A^i y))^{-R} dy \\ &\leq |\det A|^{i/2} (1 + \rho_A(x))^{-R} |\det A|^{-i(1+\theta)} \int_{\mathbb{R}^n} \rho_A(y)^\theta (1 + \rho_A(y))^{-R} dy \\ &\leq C |\det A|^{-i(1/2+\theta)} (1 + \rho_A(x))^{-R}. \end{aligned}$$

For $y \in D_2$, we have

$$\rho_A(x-y) \geq \rho_A(x)/H - \rho_A(y) \geq \rho_A(x)/H - \rho_A(x)/(2H) = \rho_A(x)/(2H),$$

and subsequently

$$(1 + \rho_A(A^i(x - y))) \geq |\det A|^i \rho_A(x - y) \geq |\det A|^i (1 + \rho_A(x)) / (4H).$$

Hence, by (2.12), (2.14), $\rho_A(y) \leq \rho_A(x) / (2H)$ and Proposition 1.3

$$\begin{aligned} \int_{D_2} &\leq \int_{D_2} \left[(1 + \rho_A(x))^{-R} + (1 + \rho_A(y))^{-R} \right] \frac{|\det A|^{i/2}}{(1 + \rho_A(A^i(x - y)))^R} dy \\ &\leq C \frac{|\det A|^{-i(R-1/2)}}{(1 + \rho_A(x))^R} \int_{\mathbb{R}^n} \frac{1}{(1 + \rho_A(y))^R} dy \leq C \frac{|\det A|^{-i(R-1/2)}}{(1 + \rho_A(x))^R} \\ &\leq C |\det A|^{-i(1/2+\theta)} (1 + \rho_A(x))^{-R}, \end{aligned}$$

since $R - 1 > \theta$.

For $y \in D_3$, we have $\rho_A(y) > \rho_A(x) / (2H)$ and, hence,

$$\begin{aligned} \int_{D_3} &\leq \int_{D_3} \left[(1 + \rho_A(x))^{-R} + (1 + \rho_A(y))^{-R} \right] \frac{|\det A|^{i/2}}{(1 + \rho_A(A^i(x - y)))^R} dy \\ &\leq C \frac{|\det A|^{i/2}}{(1 + \rho_A(x))^R} \int_{\rho_A(x-y) \geq 1} \frac{1}{\rho_A(A^i(x - y))^R} dy \\ &= C \frac{|\det A|^{-i(R-1/2)}}{(1 + \rho_A(x))^R} \int_{\rho_A(z) \geq 1} \rho_A(z)^{-R} dz \leq C |\det A|^{-i(1/2+\theta)} (1 + \rho_A(x))^{-R}. \end{aligned}$$

Combining the above estimates yields (2.16) and completes the proof of Lemma 2.3. \square

Proof of Lemma 2.1. We consider 2 cases.

Suppose that $|Q| \leq |P|$. Let $i, j \in \mathbb{Z}$ be such that $|Q| = |\det A|^{-i} \leq |\det A|^{-j} = |P|$. Then it is not hard to check that $g(x) = \Psi_P(x_P - x)$ and $h(x) = \overline{\Phi_Q(x)}$ satisfy the hypotheses of Lemma 2.3 with $R = M$, $0 < \theta < \min(M - 1, \delta)$, and $x_0 = x_Q$. Therefore,

$$\begin{aligned} |\langle \Psi_P, \Phi_Q \rangle| &= |(g * h)(x_P)| \leq C |\det A|^{-(i-j)(1/2+\theta)} (1 + |\det A|^j \rho_A(x_Q - x_P))^{-M} \\ &\leq C |\det A|^{-(i-j)(1/2+\theta)} (1 + |\det A|^j \rho_A(x_Q - x_P))^{-1-\theta}. \end{aligned}$$

Suppose that $|Q| > |P|$. Let $i, j \in \mathbb{Z}$ be such that $|Q| = |\det A|^{-j} > |\det A|^{-i} = |P|$. Again, it is not hard to check that hypotheses of Lemma 2.3 with $R = M$, $0 < \theta < \min(M - 1, \delta)$, $x_0 = x_P$, $g(x) = \overline{\Phi_Q(x_Q - x)}$, and $h(x) = \Psi_P(x)$ are satisfied. Therefore,

$$\begin{aligned} |\langle \Psi_P, \Phi_Q \rangle| &= |(g * h)(x_Q)| \leq C |\det A|^{-(i-j)(1/2+\theta)} (1 + |\det A|^j \rho_A(x_Q - x_P))^{-M} \\ &\leq C |\det A|^{-(i-j)(1/2+\theta)} (1 + |\det A|^j \rho_A(x_Q - x_P))^{-1-\theta}. \end{aligned}$$

Therefore, both cases yield

$$\begin{aligned} |\langle \Psi_P, \Phi_Q \rangle| &\leq C \left(1 + \frac{\rho_A(x_Q - x_P)}{\max(|P|, |Q|)} \right)^{-1-\theta} \cdot \begin{cases} (|Q|/|P|)^{1/2+\theta} & |Q| \leq |P| \\ (|Q|/|P|)^{1/2+\theta} & |Q| > |P| \end{cases} \\ &= C \kappa_{QP}(\theta), \end{aligned}$$

which completes the proof of Lemma 2.1. □

Theorem 2.4. *Suppose $\{\Psi_Q\}_Q \subset L^2(\mathbb{R}^n)$ is affine localized and*

$$(2.17) \quad \inf\{\|\Psi_Q\|_{L^2} : Q \in \mathcal{Q}\} > 0.$$

Then, there exists a finite partition of the collections of dilated cubes

$$\mathcal{Q} = \mathcal{Q}_1 \cup \dots \cup \mathcal{Q}_L$$

such that each of the systems $\{\Psi_Q\}_{Q \in \mathcal{Q}_i}$, $i = 1, \dots, L$ is a Riesz sequence in $L^2(\mathbb{R}^n)$. Furthermore, it is possible to choose this partition to be of the form

$$(2.18) \quad \mathcal{Q}_i = \{A^{-j}([0, 1]^n + k) : (j, k) \in (r_i + J\mathbb{Z}) \times (s_i + N\mathbb{Z}^n)\}, \quad i = 1, \dots, L,$$

where $J, N \in \mathbb{N}$, $L = JN^n$, and $(r_1, s_1), \dots, (r_L, s_L)$ are representatives of distinct cosets of $\mathbb{Z}/(L\mathbb{Z}) \times \mathbb{Z}^n/(N\mathbb{Z}^n)$.

Before we give the proof, we wish to emphasize that Theorem 2.4 goes well beyond the original idea of Gröchenig's diagonal dominance approach in [13].

Indeed, we recall that a countable set $\{\varphi_i : i \in I\}$ in a Hilbert space \mathcal{H} is a Riesz sequence if and only if the Gramian matrix G given by its entries $G_{i,j} = \langle \varphi_i, \varphi_j \rangle$ defines a bounded and invertible operator on $\ell^2(I)$. Furthermore, a sufficient condition for a matrix $(a_{i,j})_{i,j \in I}$ defining a bounded positive definite operator on $\ell^2(I)$ to be invertible is that $(a_{i,j})_{i,j \in I}$ is diagonally dominant, i.e.,

$$\inf_{i \in I} \left(|a_{i,i}| - \sum_{j \neq i} |a_{i,j}| \right) > 0.$$

The idea of Gröchenig is to decompose an index set I into a finite union of subsets $I = I_1 \cup \dots \cup I_r$, such that the corresponding Gramian matrix G is decomposed into diagonally dominant, and thus invertible on $\ell^2(I_l)$, “sub-Gramian” matrices $G^{(l)} = (G_{i,j})_{i,j \in I_l}$, where $l = 1, \dots, r$.

This approach does not work in the case of Theorem 2.4, since in Definition 2.1 the decay exponent across the scales is only $1/2 + \theta$ and one can observe that $1 + \theta$ is required for diagonal dominance approach to work. Namely, one can show that for small values $0 < \theta < 1/2$, the matrix $(\kappa_{QP})_{Q,P \in \mathcal{Q}}$ does not satisfy Schur's test of boundedness. Therefore, a more delicate approach in showing Theorem 2.4 must be used.

Proof of Theorem 2.4. Let $G = (a_{QP})_{Q,P \in \mathcal{Q}}$ be the Gramian matrix given by $a_{QP} = \langle \Psi_Q, \Psi_P \rangle$. By Lemma 2.1, we know that

$$|a_{QP}| \leq C\kappa_{QP}(\theta) \quad \text{for all } Q, P \in \mathcal{Q}.$$

Take any $J, N \in \mathbb{N}$ and for a fixed $i = 1, \dots, L = JN^n$ consider the collection of cubes $\mathcal{Q}' = \mathcal{Q}_i$ given by (2.18). Namely, for $r = 0, \dots, J - 1$ and $s \in \{0, \dots, N - 1\}^n$ consider

$$\mathcal{Q}' = \{A^{-j}([0, 1]^n + k) : (j, k) \in (r + J\mathbb{Z}) \times (s + N\mathbb{Z}^n)\}.$$

Let \mathcal{A} be the operator on $\ell^2(\mathcal{Q}')$ associated with the matrix $(a_{QP})_{Q,P \in \mathcal{Q}'}$. We decompose $\mathcal{A} = \mathcal{A}_0 + \mathcal{A}_1 + \mathcal{A}_2$ as

$$\begin{aligned} (\mathcal{A}_0 s)_Q &= a_{QQ} s_Q \\ (\mathcal{A}_1 s)_Q &= \sum_{P \in \mathcal{Q}', |P| \geq |Q|, P \neq Q} a_{QP} s_P \\ (\mathcal{A}_2 s)_Q &= \sum_{P \in \mathcal{Q}', |P| < |Q|} a_{QP} s_P, \end{aligned}$$

where $s = (s_P) \in \ell^2(\mathcal{Q}')$.

It is clear that \mathcal{A}_0 is an invertible diagonal operator on $\ell^2(\mathcal{Q}')$, since

$$\inf\{a_{QQ} : Q \in \mathcal{Q}'\} \geq \inf\{\|\Psi_Q\|_{L^2}^2 : Q \in \mathcal{Q}\} > 0.$$

Our goal is to show that for appropriate choice of $J, N \in \mathbb{N}$, \mathcal{A}_1 and \mathcal{A}_2 have very small operator norms on $\ell^2(\mathcal{Q}_i)$ for every $i = 1, \dots, JN^n$. Therefore, \mathcal{A} must be also invertible on $\ell^2(\mathcal{Q}_i)$ as a small perturbation of an invertible operator \mathcal{A}_0 .

Indeed, take any $\varepsilon > 0$ and let $s = (s_Q) \in \mathcal{Q}'$. By the Cauchy-Schwartz inequality

$$\begin{aligned} (2.19) \quad & \sum_{Q \in \mathcal{Q}'} |(\mathcal{A}_1 s)_Q|^2 \\ & \leq C \sum_{Q \in \mathcal{Q}'} \left| \sum_{P \in \mathcal{Q}', |P| \geq |Q|, P \neq Q} (1 + \rho_A(x_P - x_Q)/|P|)^{-1-\theta} \left(\frac{|Q|}{|P|}\right)^{1/2+\theta} |s_P| \right|^2 \\ & \leq C \sum_{Q \in \mathcal{Q}'} \left[\sum_{P \in \mathcal{Q}', |P| \geq |Q|, P \neq Q} (1 + \rho_A(x_P - x_Q)/|P|)^{-1-\theta} \left(\frac{|Q|}{|P|}\right)^\theta \right] \\ & \quad \times \left[\sum_{P \in \mathcal{Q}', |P| \geq |Q|, P \neq Q} (1 + \rho_A(x_P - x_Q)/|P|)^{-1-\theta} \left(\frac{|Q|}{|P|}\right)^{1+\theta} |s_P|^2 \right] \end{aligned}$$

Note that the expression in the first bracket is finite, since

$$\sum_{P \in \mathcal{Q}, |P| = |\det A|^{-j}} (1 + \rho_A(x_P - x_Q)/|P|)^{-1-\theta} \leq C < \infty,$$

independent of $j \in \mathbb{Z}$. Therefore, by (2.19) and Lemma 2.2

$$\begin{aligned} (2.20) \quad & \sum_{Q \in \mathcal{Q}'} |(\mathcal{A}_1 s)_Q|^2 \leq C \sum_{P \in \mathcal{Q}'} |s_P|^2 I_P, \quad \text{where} \\ & I_P = \sum_{Q \in \mathcal{Q}', |Q| \leq |P|, Q \neq P} (1 + \rho_A(x_P - x_Q)/|P|)^{-1-\theta} \left(\frac{|Q|}{|P|}\right)^{1+\theta} \end{aligned}$$

To estimate I_P , we pick any $P = A^{-r-Jj_0}([0, 1]^n + s + Nk_0) \in \mathcal{Q}'$, where $j_0 \in \mathbb{Z}$, $k_0 \in \mathbb{Z}^n$. Next, we split the summation over $|Q| = |P|$ and $|Q| < |P|$

$$\begin{aligned} & \sum_{Q \in \mathcal{Q}', |Q|=|P|, Q \neq P} (1 + \rho_A(x_P - x_Q)/|P|)^{-1-\theta} \left(\frac{|Q|}{|P|} \right)^{1+\theta} \\ &= \sum_{k \in \mathbb{Z}^n, k \neq k_0} (1 + \rho_A(Nk_0 - Nk))^{-1-\theta} = \sum_{k \in \mathbb{Z}^n \setminus \{0\}} (1 + \rho_A(Nk))^{-1-\theta} < \varepsilon \end{aligned}$$

for sufficiently large N . Likewise, by Lemma 2.2

$$\begin{aligned} & \sum_{Q \in \mathcal{Q}', |Q| < |P|} (1 + \rho_A(x_P - x_Q)/|P|)^{-1-\theta} \left(\frac{|Q|}{|P|} \right)^{1+\theta} \\ &= \sum_{j=j_0+1}^{\infty} \sum_{k \in \mathbb{Z}^n} \frac{|\det A|^{J(j_0-j)(1+\theta)}}{(1 + |\det A|^{r+Jj_0} \rho_A(A^{-r-Jj}(s + Nk) - x_P))^{1+\theta}} \\ &\leq \sum_{j=j_0+1}^{\infty} \sum_{k \in \mathbb{Z}^n} \frac{|\det A|^{J(j_0-j)(1+\theta)}}{(1 + \rho_A(A^{J(j_0-j)}(s + k) - A^{r+Jj_0}x_P))^{1+\theta}} \\ &\leq C \sum_{j=j_0+1}^{\infty} |\det A|^{J(j_0-j)\theta} = C |\det A|^{-J} / (1 - |\det A|^{-J}) < \varepsilon \end{aligned}$$

for sufficiently large J . Combining the above estimates, which are independent of $P \in \mathcal{Q}'$, with (2.20) yields

$$\|\mathcal{A}_1 s\|_{\ell^2(\mathcal{Q}')}^2 \leq 2\varepsilon \|s\|_{\ell^2(\mathcal{Q}')}^2.$$

Analogously, starting from the estimate

$$\begin{aligned} \sum_{Q \in \mathcal{Q}'} |(\mathcal{A}_2 s)_Q|^2 &\leq C \sum_{Q \in \mathcal{Q}'} \left[\sum_{P \in \mathcal{Q}', |P| < |Q|} (1 + \rho_A(x_P - x_Q)/|Q|)^{-1-\theta} \left(\frac{|Q|}{|P|} \right)^{1+\theta} \right] \\ &\quad \times \left[\sum_{P \in \mathcal{Q}', |P| > |Q|} (1 + \rho_A(x_P - x_Q)/|Q|)^{-1-\theta} \left(\frac{|Q|}{|P|} \right)^\theta |s_P|^2 \right], \end{aligned}$$

and using similar arguments as above one can obtain

$$\|\mathcal{A}_2 s\|_{\ell^2(\mathcal{Q}')}^2 \leq 2\varepsilon \|s\|_{\ell^2(\mathcal{Q}')}^2.$$

Since $\varepsilon > 0$ is arbitrary, this shows that \mathcal{A}_1 and \mathcal{A}_2 have small operator norms and consequently \mathcal{A} is invertible on $\ell^2(\mathcal{Q}')$ for sufficiently large $J, N \in \mathbb{N}$. This completes the proof of Theorem 2.4. \square

Corollary 2.5 follows immediately from Theorem 2.4 and Remark 2.2.

Corollary 2.5. *Suppose $\psi(x)$ is a non-zero function such that the $\{\psi_Q : Q \in \mathcal{Q}\}$ is affine localized. In particular, it suffices to assume that $\psi(x)$ satisfies*

$$\begin{aligned} |\psi(x) - \psi(y)| &\leq C|x - y|^\delta(1 + |x|)^{-M/\zeta^-} && \text{for } |x - y| \leq 1, \\ |\psi(x)| &\leq C(1 + |x|)^{-M/\zeta^-}, \\ \int_{\mathbb{R}^n} \psi(x)dx &= 0, \end{aligned}$$

for some $M > 1$, $\delta > 0$, and $C > 0$.

Then $\{\psi_Q : Q \in \mathcal{Q}\}$ is a finite union of pairwise disjoint affine systems, each of which is a Riesz sequence. More precisely, there exist $J, N \in \mathbb{N}$ such that the affine systems

$$(2.21) \quad \{D^j T_k(D^{r_i} T_{s_i} \psi) : j \in J\mathbb{Z}, k \in A^{-r_i} N\mathbb{Z}^n\}, \quad i = 1, \dots, JN^n$$

form a partition of $\{\psi_Q : Q \in \mathcal{Q}\}$ and each system in (2.21) is a Riesz sequence, where (r_i, s_i) 's are the same as in Theorem 2.4.

We remark that in Corollary 2.5 it is not assumed that $\psi(x)$ is a framelet. Nonetheless, the case when $\psi(x)$ is a framelet is probably the most interesting, since it provides a partial answer to the Feichtinger conjecture. In particular, it shows that if a frame wavelet generates affine system $\{\psi_Q : Q \in \mathcal{Q}\}$ localized in the sense of Definition 2.1, then it can be split into a finite union of systems each of which is a Riesz sequence. Since each of these systems is again an affine system (with respect to some power of dilation A and lattice given by (2.21)), a stronger variant of the Feichtinger conjecture holds under the localization assumption.

We also wish to emphasize that Corollary 2.5 is optimal, at least in the qualitative sense. That is, if we remove the hypothesis of $\{\psi_Q : Q \in \mathcal{Q}\}$ being affine localized then the conclusion of Corollary 2.5 is in general invalid even if $\psi(x)$ is a framelet. Therefore, we will show that the stronger variant of the Feichtinger conjecture for wavelet frames, asserting that affine frame systems can be split into a finite disjoint union of affine Riesz sequences, does not hold without a localization assumption on $\psi(x)$.

Theorem 2.6. *There exists a tight frame wavelet $\psi(x) \in L^2(\mathbb{R})$ such that the stronger variant of the Feichtinger conjecture fails. More precisely, $\{D^{Jj} T_{Nk} \psi : j, k \in \mathbb{Z}\}$ is never a Riesz sequence for any choice of $J, N \in \mathbb{N}$. Moreover, $\psi(x)$ can be chosen to be a semi-orthogonal frame wavelet.*

Proof. Assume momentarily that we find a nowhere dense set $W \subset [-1/2, 1/2]$ such that $\{2^j W : j \in \mathbb{Z}\}$ partitions \mathbb{R} (modulo null sets) and the sets $\{W + k : k \in \mathbb{Z}\}$ are pairwise disjoint (modulo null sets). Later, we will show how to construct such W .

Let $\psi \in L^2(\mathbb{R})$ be given by $\hat{\psi} = \chi_W$. A standard support argument shows that $D^j T_k \psi \perp D^{j'} T_{k'} \psi$ for all $j \neq j' \in \mathbb{Z}$ and $k, k' \in \mathbb{Z}$. Moreover, since

$$\sum_{k \in \mathbb{Z}} |\hat{\psi}(\xi + k)|^2 = \chi_{\tilde{W}}(\xi), \quad \text{where } \tilde{W} = \bigcup_{k \in \mathbb{Z}} (k + W),$$

$\{T_k\psi : k \in \mathbb{Z}\}$ is a tight frame with constant 1 for its closed linear span, which consists of all $f \in L^2(\mathbb{R}^n)$ with $\text{supp } \hat{f} \subset W$. Hence, $\{D^j T_k \psi : k \in \mathbb{Z}\}$ has the same property for any $j \in \mathbb{Z}$. Since $\{2^j W : j \in \mathbb{Z}\}$ partitions \mathbb{R} , ψ is a semi-orthogonal tight frame wavelet. Moreover, since $W \subset [-1/2, 1/2]$ is nowhere dense we have that $\sum_{k \in \mathbb{Z}} |\hat{\psi}(\xi + k/N)|^2$ must vanish on a set of positive measure and hence $\{T_{kN}\psi : k \in \mathbb{Z}\}$ is not a Riesz sequence for any $N \in \mathbb{N}$. Consequently, $\{D^{Jj} T_{Nk}\psi : j, k \in \mathbb{Z}\}$ is never a Riesz sequence for any choice of $J, N \in \mathbb{N}$.

Hence, it remains to show the existence of a set W . Let $\{W_k : k = 1, 2, \dots\} \subset [-1, -1/2] \cup [1/2, 1]$ be a family of closed nowhere dense sets such that

$$\bigcup_{k=1}^{\infty} W_k = [-1, -1/2] \cup [1/2, 1] \quad \text{and} \quad \sum_{k=1}^{\infty} |W_k| = 1.$$

Clearly, W_k 's must be pairwise disjoint modulo null sets. Define $W = \bigcup_{k=1}^{\infty} 2^{-k} W_k$. An easy verification shows that W satisfies all required properties. \square

3. GABOR FRAMES

As mentioned in Section 2, Gröchenig in [13] introduced a notion of intrinsically localized frames that are well suited for systems (such as Gabor systems) which are generated by families of unitary operators with compatible actions on $L^2(\mathbb{R}^n)$. In particular, he was able to show that if g is in a certain modulation space, then the (possibly irregular) Gabor system $\{M^k T^l g : (k, l) \in X\}$ can be written as the finite union of Riesz basic sequences provided X is the finite union of separated sets. In particular, the Feichtinger conjecture holds for such Gabor systems.

An earlier result in this direction was provided by Ron and Shen [21], which we describe now. Let $K, L \subset \mathbb{R}^n$ be full rank lattices and $K^*, L^* \subset \mathbb{R}^n$ be their dual lattices. The Gabor system $\{M^k T^l g : k \in K, l \in L\}$ is said to be *sup-adjoint* if $K^* \subset L$. Ron and Shen showed in [21, Theorem 3.5] that if $\{M^k T^l g : k \in K, l \in L\}$ is sup-adjoint and a frame for $L^2(\mathbb{R}^n)$, then it can be written as the finite union of Riesz basic sequences. Note that in the one-dimensional case, this corresponds to $\{M^{ma} T^{nb} g : m, n \in \mathbb{Z}\}$ with $ab = 1/N$ for integer $N \geq 1$.

Sticking to regular systems, Casazza et al. [6, Theorem 5.1] were able to generalize the one-dimensional version of Ron and Shen's theorem by showing that if the Gabor system $\{M^{ma} T^{nb} g : m, n \in \mathbb{Z}\}$ is a frame for $L^2(\mathbb{R}^n)$ and ab is rational, then the Gabor system can be written as the finite union of Riesz basic sequences.

Ron and Shen proposed [21] a class of Gabor systems that are higher dimensional analogs of rational Gabor systems, which we describe here.

Definition 3.1. [21] Let K, L be full rank lattices in \mathbb{R}^n . The Gabor system $\{M^k T^l g : k \in K, l \in L\}$ is said to be *compressible* if $L^* \cap K$ has rank n .

Remark 3.1. In the one dimensional case $\{M^{ma} T^{nb} g : m, n \in \mathbb{Z}\}$ is a compressible Gabor system if and only if ab is rational.

The main result of this short section is the following, which is a generalization of both [21, Theorem 3.5] and [6, Theorem 5.1].

Theorem 3.1. *Let $K, L \subset \mathbb{R}^n$ be full-rank lattices, and $g \in L^2(\mathbb{R}^n)$. If $\{M^k T^l g : k \in K, l \in L\}$ is a compressible Gabor system which is a frame for $L^2(\mathbb{R}^n)$, then it can be written as the finite union of Riesz basic sequences.*

Before proving this theorem, we will develop some tools. The following simple proposition is the key to our development.

Proposition 3.2. *Let $K, L \subset \mathbb{R}^n$ be full-rank lattices such that $K^* \cap L$ has rank n as well. Then, there exist $I < \infty$ and $\{a_i : i = 1, \dots, I\}$ such that $\Gamma := \cup_{i=1}^I (L + a_i)$ is a lattice, and $K^* \cup L \subset \Gamma$.*

Proof. Choose $\{a_i : i = 1, \dots, I\} \subset K^*$ to be a complete set of representatives of $K^*/(K^* \cap L)$. Then, $K^* = \cup_{i=1}^I ((K^* \cap L) + a_i)$. We claim $\Gamma = \cup_{i=1}^I (L + a_i)$ satisfies the conclusion of the theorem. Indeed, $K^* \subset \Gamma$ follows immediately from the definition of Γ , and $L \subset \Gamma$ follows from the fact that one a_i is in L .

To show Γ is a lattice, it suffices to show that Γ is a group, since it is clearly a discrete subset of \mathbb{R}^n . Let $x_1 = l_1 + a_1$ and $x_2 = l_2 + a_2$ be in Γ , where a_1, a_2 are intended to represent arbitrary elements of $\{a_i : i = 1, \dots, I\}$, and l_j are arbitrary elements of L . Then, since $K^*/(K^* \cap L)$ is a group, there is a number $k \in \{1, \dots, I\}$ such that $(K^* \cap L + a_1) - (K^* \cap L + a_2) = (K^* \cap L + a_k)$, i.e., $a_k - (a_1 - a_2) = l \in K^* \cap L$. Then, $x_1 - x_2 = l_1 - l_2 - l + a_k \in L + a_k \subset \Gamma$. \square

Remark 3.2. In the one-dimensional case, if $\{M^k T^l g : k \in \mathbb{Z}, l \in (p/q)\mathbb{Z}\}$ with $0 < p/q < 1$ is a compressible Gabor system, then Proposition 3.2 yields $\Gamma = \cup_{i=0}^{p-1} (p/q\mathbb{Z} + i)$. Casazza et al. in their proof that one-dimensional compressible Gabor systems satisfy the Feichtinger conjecture also found this oversampling, but wrote it as $\Gamma = \cup_{i=0}^{p-1} (p/q\mathbb{Z} + i/q)$.

We now show that the condition on the lattices in Proposition 3.2 is equivalent to the Gabor system being compressible.

Proposition 3.3. *Let $K, L \subset \mathbb{R}^n$ be full-rank lattices. Then, $K^* \cap L$ is rank n if and only if $K \cap L^*$ is rank n .*

Proof. Suppose $K^* \cap L$ has rank n . Obtain Γ from Proposition 3.2. Since $L \cup K^* \subset \Gamma$, it follows that $\Gamma^* \subset L^* \cap K$. Hence, since Γ^* has rank n , so does $L^* \cap K$. \square

We now prove the main result in this section.

Proof of Theorem 3.1. Let Γ be as in Proposition 3.2. Then, $\{M^k T^\gamma g : k \in K, \gamma \in \Gamma\}$ is a sup-adjoint Gabor system. Since $L \subset \Gamma$, a lower frame bound is satisfied, and since the cardinality of I is finite, and $\Gamma = \cup_{i \in I} (L + a_i)$ an upper frame bound is also satisfied. By [21, Theorem 3.5], $\{M^k T^\gamma g : k \in K, \gamma \in \Gamma\}$ can be written as the finite union of Riesz basic sequences. Hence, since $\{M^k T^l g : k \in K, l \in L\} \subset \{M^k T^\gamma g : k \in K, \gamma \in \Gamma\}$, the result holds. \square

4. FRAMES OF TRANSLATES

In this section we investigate the following question. Suppose that

$$\mathcal{S}_f^\Gamma := \{f(x + k) : k \in \Gamma\}, \quad \text{where } \Gamma \subset \mathbb{R}^n$$

is a frame for its closed linear span. Can \mathcal{S}_f^Γ be written as the finite union of Riesz basic sequences? There is a large collection of results that can be applied to this question. One purpose of this section is to show what types of results can be proven using known techniques, and to show where the limits of known techniques seem to be. Of particular interest is Theorem 4.13, which provides evidence against the Feichtinger conjecture even in this restricted setting.

We begin by applying the results of Gröchenig [13] to this setting. Let \mathcal{X} be a *separated* subset of \mathbb{R}^n , i.e., $\inf_{x,y \in \mathcal{X}, x \neq y} |x - y| = \delta > 0$. Recall that a collection of vectors $\{e_x : x \in \mathcal{X}\} \subset \mathcal{H}$ is said to be *intrinsically localized* if for some $s > n$,

$$|\langle e_x, e_y \rangle| \leq C(1 + |x - y|)^{-s}$$

for all $x, y \in \mathcal{X}$.

Proposition 4.1. *Let $f \in L^2(\mathbb{R}^n)$ and Γ be a separated subset of \mathbb{R}^n . If $|f(x)| \leq (1 + |x|)^{-s}$ for some $s > n$, then \mathcal{S}_f^Γ is intrinsically localized. In particular, if Γ is the finite union of separated subsets of \mathbb{R}^n and \mathcal{S}_f^Γ is a frame for its span, then it can be written as the finite union of Riesz basic sequences.*

Proof. It suffices to show for $k, l \in \Gamma$ that

$$|\langle f(x+k), f(x+l) \rangle| = |\langle f(x), f(x+(l-k)) \rangle| \leq C(1 + |l-k|)^{-s}.$$

Write $m = l - k$. Let $I_1 = \{x \in \mathbb{R}^n : |x| < |x+m|\}$, and $I_2 = \mathbb{R}^n \setminus I_1$. Then, for $x \in I_1$, $|x+m| \geq \frac{1}{2}|m|$, and for $x \in I_2$, $|x| \geq \frac{1}{2}|m|$. Thus,

$$\begin{aligned} |\langle f(x), f(x+m) \rangle| &= \left| \int_{I_1} f(x) \overline{f(x+m)} dx + \int_{I_2} f(x) \overline{f(x+m)} dx \right| \\ &\leq \int_{I_1} |f(x)| |f(x+m)| dx + \int_{I_2} |f(x)| |f(x+m)| dx \\ &\leq (1 + |m/2|)^{-s} (2 \int |f(x)| dx), \end{aligned}$$

as desired.

For the second statement, since $\Gamma = \cup_{i=1}^M \Gamma_i$, with each Γ_i separated, one can apply [13, Theorem 2] to each Γ_i separately and the result follows. \square

Remark 4.1. By analyzing the proof of [13, Theorem 2] more carefully (see also Theorem 2.4 above), in the case that Γ is a lattice and \mathcal{S}_f^Γ is intrinsically localized, one can show that the Riesz basic sequences can all be chosen to be translates of a function by a lattice. Thus, the stronger version of the Feichtinger conjecture holds in this case. We omit the details.

In the case that Γ is a lattice, one can weaken the hypotheses of Proposition 4.1 somewhat. Denote

$$(4.1) \quad \sigma_f^\Gamma(\xi) = \sum_{\gamma \in \Gamma^*} |f(\xi + \gamma)|^2,$$

where we will suppress the dependence on Γ when it is obvious from context. We begin by recalling the following, see [5].

Theorem 4.2. *Let $f \in L^2(\mathbb{R}^n)$, and $\Gamma \subset \mathbb{R}^n$ a full-rank lattice. Then,*
(i) \mathcal{S}_f^Γ is a frame for its closed linear span if and only if there exists $0 < A \leq B < \infty$ such that $A\chi_U \leq \sigma_f^\Gamma \leq B\chi_U$ almost everywhere, where $U = \{\xi \in \mathbb{R}^n : \sigma_f^\Gamma(\xi) > 0\}$.
(ii) \mathcal{S}_f^Γ is a Riesz basic sequence if and only if there exists $0 < A \leq B < \infty$ such that $A \leq \sigma_f^\Gamma \leq B$ almost everywhere.

An immediate application of Theorem 4.2 yields

Proposition 4.3. *Let $f \in L^2(\mathbb{R}^n)$, $\Gamma \subset \mathbb{R}^n$ a full-rank lattice. If \mathcal{S}_f^Γ is a frame for its closed linear span and there exists $\eta = (\eta_1, \dots, \eta_n) \in [0, 1]^n$, $\epsilon > 0$ such that*

$$(4.2) \quad E := \eta + B_\epsilon(0) \subset \text{supp}(\sigma_f^\Gamma),$$

then \mathcal{S}_f^Γ can be written as the finite union of Riesz basic sequences.

Proof. Choose a lattice Λ of rank n satisfying $\Gamma^* \subset \Lambda$ and

$$(4.3) \quad \sum_{\lambda \in \Lambda} \chi_E(\xi + \lambda) \geq 1, \quad \text{a.e.}$$

Then, $A \leq \sigma_f^\Lambda \leq B$ almost everywhere, so $\mathcal{S}_f^{\Lambda^*}$ is a Riesz basic sequence. Since $\Gamma = \cup_{i \in I} (\Lambda^* + a_i)$, and $\mathcal{S}_f^{\Lambda^* + a_i}$ are all Riesz basic sequences, the proposition follows. \square

Remark 4.2. Proposition 4.3 applies, in particular, if $\hat{f} = \chi_{[a,b]}$. In this case,

$$|\langle f, T^k f \rangle| = \frac{1}{2\pi|k|} |e^{2\pi i k b} - e^{2\pi i k a}|,$$

so for almost every choice of a, b , f is not intrinsically localized. However, for every choice of a and b , $\mathcal{S}_f^{\mathbb{Z}}$ is a frame and can be written as the finite union of Riesz basic sequences.

With a little more work, we can improve on Propositions 4.3 and 4.1 in the one dimensional case. Recall the following theorem of Montgomery and Vaughan [19, Theorem 1, Chapter 7], [20].

Theorem 4.4. *Suppose that $\lambda_1, \dots, \lambda_N$ are distinct real numbers, and suppose that $\delta > 0$ is chosen so that $|\lambda_n - \lambda_m| \geq \delta$ whenever $n \neq m$. Then, for any coefficients a_1, \dots, a_N , and any $T > 0$,*

$$(4.4) \quad (T - 1/\delta) \sum_{n=1}^N |a_n|^2 \leq \int_0^T \left| \sum_{n=1}^N a_n e^{2\pi i \lambda_n t} \right|^2 dt \leq (T + 1/\delta) \sum_{n=1}^N |a_n|^2.$$

We will also need the following lemma, which has several variants in the literature, e.g. [16, Lemma 1]. We include a proof for completeness.

Lemma 4.5. *Let $f \in L^2(\mathbb{R})$ and $\Lambda = \{\lambda_n : n \in \mathbb{Z}\}$. If there exists a constant B such that for every $m \in \mathbb{Z}$*

$$\sum_{n \in \mathbb{Z}} |\langle e^{2\pi i \lambda_n t} f, e^{2\pi i \lambda_m t} f \rangle|^2 \leq B,$$

then the number of points λ_n inside each interval of length 1 is uniformly bounded.

Proof. We prove the contrapositive. By standard arguments, there exists an $\epsilon > 0$ such that for all η with $|\eta| \leq \epsilon$,

$$(4.5) \quad \left| \int_{\mathbb{R}} e^{2\pi i \eta t} |f(t)|^2 dt \right|^2 \geq \frac{1}{2} \|f\|^2.$$

Suppose that the number of λ_n inside an interval of length one is not bounded. Then, the number of λ_n inside intervals of length ϵ is also not bounded. Choose μ_k such that the number of λ_n inside $A_k := [\mu_k - \epsilon/2, \mu_k + \epsilon/2]$ is at least k . Choose $\lambda_{n_k} \in A_k \cap \Lambda$ arbitrarily. Then, by (4.5) it follows that

$$\sum_{n \in \mathbb{Z}} |\langle e^{2\pi i \lambda_{n_k} t} f, e^{2\pi i \lambda_n t} f \rangle|^2 \geq \frac{k}{2} \|f\|^2.$$

Hence, Λ does not satisfy the hypotheses of the lemma. \square

Theorem 4.6. *Let $f \in L^2(\mathbb{R})$ be such that there exists $a > 0$ and an interval I such that $a\chi_I \leq \hat{f}$, and let Γ be an arbitrary subset of \mathbb{R} . If S_f^Γ is a frame for its span, then S_f^Γ can be written as the finite union of Riesz basic sequences.*

Proof. By Parseval's identity, $\langle f(x + \gamma_1), f(x + \gamma_2) \rangle = \langle e^{2\pi i \gamma_1 \xi} \hat{f}(\xi), e^{2\pi i \gamma_2} \hat{f}(\xi) \rangle$. Thus, since S_f^Γ is a frame in its span, the hypotheses of Lemma 4.5 are satisfied, and Γ can be written as the finite union of separated sequences. In particular, one can write Γ as the finite union of sets $\Gamma = \Gamma_1 \cup \dots \cup \Gamma_N$ such that for each $1 \leq i \leq N$, $\delta_i := \inf\{|\gamma - \lambda| : \gamma, \lambda \in \Gamma_i, \gamma \neq \lambda\}$ satisfies $\frac{1}{\delta_i} < |I|$. Clearly, the interval $[0, T]$ in Theorem 4.4 can be replaced by any interval of length T , so $\{f(x + \gamma) : \gamma \in \Gamma_i\}$ is a Riesz basic sequence with bounds $|I| \pm \frac{1}{\delta_i}$. \square

Corollary 4.7. *If $\{e^{2\pi i \lambda \xi} : \lambda \in \Lambda\}$ is a Fourier frame, i.e., a frame for $L^2([0, 1])$, then it can be written as the finite union of Riesz basic sequences.*

Corollary 4.7 was also observed by Christensen and Lindner [8], see the discussion after Theorem 2.7 of their paper.

4.1. Frames of integer translates. For the rest of this section we will restrict our attention to the special case of frames of translates S_f^Γ with $\Gamma = \mathbb{Z}$. In this setting, we can show that the following relationship between frames of translates and pavings of Laurent operators.

Proposition 4.8. *Let $f \in L^\infty(\mathbb{T})$ be such that $S_f^\mathbb{Z}$ is a frame for its span and \hat{f} is a bounded function supported on \mathbb{T} . If the Laurent operator $L_{|\hat{f}|^2}$ satisfies the Paving conjecture, then $S_f^\mathbb{Z}$ can be written as the finite union of Riesz basic sequences.*

Proof. According to [6, Proposition 3.1] and its proof, a frame $\{f_n\}$ can be written as the finite union of Riesz basic sequences if the operator S , whose matrix is given by

$$\langle S e_i, e_j \rangle = \begin{cases} \langle f_i, f_j \rangle & i \neq j \\ 0 & i = j \end{cases}$$

can be paved. Applying this to the frame $\{e^{2\pi ik\xi}\hat{f}(\xi) : k \in \mathbb{Z}\}$ yields S of the form

$$\langle Se_i, e_j \rangle = \begin{cases} |\hat{f}|^2(i-j) & i \neq j \\ 0 & i = j \end{cases}.$$

This is exactly the matrix of $L_{|\hat{f}|^2}$ minus its diagonal; hence, the result follows. \square

Theorem 4.6 should be contrasted with the following elementary observation.

Example 4.9. *Let U be a nowhere dense subset of $[0, 1]$ with positive measure, $\hat{f} = \chi_U$. Then, since $[0, 1]$ cannot be written a.e. as the finite union of nowhere dense sets, for any natural number $N > 0$,*

$$(4.6) \quad \sum_{k \in \mathbb{Z}} \chi_U(\xi + k/N) = 0$$

on a set of positive measure. Hence, by Theorem 4.2, $\mathcal{S}_f^{N\mathbb{Z}}$ is not a Riesz basic sequence.

While there is a characterization of when $\{f(x+k) : k \in \mathcal{K}\}$ is a Riesz basic sequence for $\mathcal{K} \subset \mathbb{Z}$ [5], it is still difficult to decide whether $\{f(x+k) : k \in \mathcal{K}\}$ is a Riesz basic sequence when Γ is an irregular subset of \mathbb{Z} and f is poorly localized. In particular, for f as generally defined as in Example 4.9, the authors do not know whether the Feichtinger conjecture is satisfied. We formalize the most natural question along this line of thought as:

Question 4.10. *Suppose $\hat{f} = \chi_U$, and $\{f(x+k) : k \in \mathbb{Z}\}$ is a frame for its closed linear span. Can $\{f(x+k) : k \in \mathbb{Z}\}$ be written as the finite union of Riesz basic sequences?*

Of course, by Proposition 4.8, a positive solution to the paving problem for Laurent operators would imply a positive answer to Question 4.10.

We present now a specially chosen f of the form $\hat{f} = \chi_U$ with the property that any set $\mathcal{K} \subset \mathbb{Z}$ for which $\{f(x+k) : k \in \mathcal{K}\}$ is a Riesz basic sequence must be “quite strange”. While this example does not answer Question 4.10, it will suggest a line of attack for studying it. We will need to recall some machinery in order to define $\hat{f} = \chi_U$. We follow the development given in [14].

Let $u = \{u_n : n = 1, 2, \dots\}$ be a sequence of points in the torus $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. The sequence u is said to be *uniformly distributed* if

$$\lim_{N \rightarrow \infty} \frac{\#\{1 \leq n \leq N : u_n \in I\}}{N} = |I|$$

for every arc I in the torus. For a finite, nonempty set $S \subset \mathbb{T}$ (possibly with multiplicity), we define the *discrepancy* of S to be

$$\text{Discr}(S) = \sup_{I \subset \mathbb{T}} \left| \frac{\#(S \cap I)}{\#S} - |I| \right|.$$

The N -term discrepancy of the sequence u is defined as

$$D_N(u) = \text{Discr}(\{u_n : 1 \leq n \leq N\}).$$

We will need two results.

Theorem 4.11. [19, Koksma's Inequality] *For any sequence of points u_1, \dots, u_N in \mathbb{T} , and any function $f : \mathbb{T} \rightarrow \mathbb{R}$ of bounded variation,*

$$\left| \frac{1}{N} \sum_{n=1}^N f(u_n) - \int_0^1 f(t) dt \right| \leq \text{Var}(f) \text{Discr}(\{u_n : 1 \leq n \leq N\}),$$

where $\text{Var}(f)$ is the total variation of f .

Theorem 4.12. [17] *For any $\epsilon > 0$, the N -term discrepancy of $u_n = n\alpha \pmod{1}$ satisfies*

$$D_N(u) \leq C_\alpha N^{-1} \log^{2+\epsilon} N$$

for almost all α , where C_α depends only on α .

Definition 4.1. Let $g : \mathbb{R}^2 \rightarrow \mathbb{R}$. We say that $\mathcal{K} \subset \mathbb{Z}$ satisfies the $g(\ell, N)$ arithmetic progression condition if for every $\delta > 0$ there exists $M, N, \ell \in \mathbb{Z}$ such that

- (i) $g(\ell, N) < \delta$, and
- (ii) $\{M, M + \ell, \dots, M + N\ell\} \subset \mathcal{K}$.

Theorem 4.13. *There exists a set $U \subset [0, 1]$ such that if $\mathcal{K} \subset \mathbb{Z}$ satisfies the $g(\ell, N) = \ell N^{-1/2} \log^3 N$ arithmetic progression condition, then $\{f(x + k) : k \in \mathcal{K}\}$ is not a Riesz basic sequence, where $\hat{f} = \chi_U$.*

Proof. Let $u = u_N(\xi) = \{\xi, 2\xi, \dots, N\xi\}$. By Theorem 4.12 with $\epsilon = 1$, for almost all ξ and all N ,

$$D_N(u) \leq \frac{C(\xi)}{N} \log^3 N.$$

Choose K such that

$$0 < |\{\xi \in [0, 1] : C(\xi) \leq K\}| < 1,$$

and let $U = \{\xi \in [0, 1] : C(\xi) \leq K\}$. (This choice of K is possible since C is not a constant function. However, it formally doesn't matter for this proof, since if C were constant, then one could choose any subset of $[0, 1]$.)

Let $\delta > 0$ be arbitrary and find M, N and ℓ such that $\ell N^{-1/2} \log^3 N < \delta$ and such that $\{M, M + \ell, \dots, M + \ell N\} \subset \mathcal{K}$. Let $f(u) = e^{2\pi i \ell u}$. By Theorem 4.11,

$$\left| \frac{1}{N} \sum_{k=1}^N f(k\xi) - 0 \right| \leq \text{Var}(f) \text{Discr}(u_N(\xi)).$$

Therefore,

$$\left| \frac{1}{N} e^{2\pi i M \xi} \sum_{k=1}^N f(k\xi) \right| \leq K \ell \frac{1}{N} \log^3 N.$$

Multiplying both sides by \sqrt{N} and integrating yields

$$\left(\int_U \left| \frac{1}{\sqrt{N}} e^{2\pi i M \xi} \sum_{k=1}^N f(k\xi) \right|^2 \right)^{1/2} \leq |U|^{1/2} K \ell N^{-1/2} \log^3 N < \delta.$$

So, since $\sum_{k=1}^N \frac{1}{\sqrt{N}}^2 = 1$, it follows from the definition of Riesz basis that $\{f(x+k) : k \in \mathcal{K}\}$ is not a Riesz basic sequence. \square

Remark 4.3. Theorem 4.13 should be compared to [15, Theorem 5.4 (b)], where a similar negative result is shown for pavings of Laurent operators of certain types. Since the Paving Conjecture for Laurent operators implies the Feichtinger conjecture for frames of translations (Proposition 4.8), the example in Theorem 4.13 is a stronger negative example. The examples given in [15, Theorem 5.4 (b)] were considered good candidates for being counterexamples to the Paving Conjecture until it was proven in [2] that many such operators can be paved.

Remark 4.4. If one randomly assigns each integer to one of L subsets $\mathcal{K}_1, \dots, \mathcal{K}_L$, then with probability one, for each i and L there will exist M_i such that

$$\{M_i, M_i + 1, \dots, M_i + L\} \subset \mathcal{K}_i.$$

Hence, with f as in Theorem 4.13 and \mathbb{Z} partitioned as $\cup_{i=1}^L \mathcal{K}_i$, the probability that the set $\{f(x+k) : k \in \mathcal{K}_i\}$ is a Riesz basic sequence is zero.

One should also compare the $\ell N^{-1/2} \log^3 N$ arithmetic progression condition with van der Waerden's theorem, which states that, given a partition of the integers $\{\mathcal{K}_1, \dots, \mathcal{K}_L\}$, there exists an i such that \mathcal{K}_i has arbitrarily long arithmetic progressions. One might hope that there is a quantitative version of van der Waerden's theorem that would give the existence of $\ell N^{-1/2} \log^3 N$ arithmetic progressions, but this is not the case. The current best quantitative version is given by Gowers [12], where we use the notation $a \uparrow b$ to denote a^b , with natural rules for order of operation.

Theorem 4.14. [12, Theorem 18.2] *Let $0 < \gamma \leq 1/2$, let k be a positive integer, let $P \geq 2 \uparrow 2 \uparrow \gamma^{-1} \uparrow 2 \uparrow 2 \uparrow (k+9)$, and let A be a subset of $\{1, \dots, P\}$ of size at least γP . Then A contains an arithmetic progression of length k .*

Corollary 4.15. *Suppose \mathbb{Z} is partitioned into L subsets $\mathcal{K}_1, \dots, \mathcal{K}_L$. Let $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by $g(\ell, N) = \frac{\ell}{2 \uparrow 2 \uparrow L \uparrow 2 \uparrow 2 \uparrow (N+9)}$. Then, there exists an i such that the set \mathcal{K}_i satisfies the $g(\ell, N)$ arithmetic progression condition.*

Proof. With g as above, let $\delta = \frac{1}{N}$. Choose $\gamma = \frac{1}{L}$, $k = N$ and $P = 2 \uparrow 2 \uparrow L \uparrow 2 \uparrow 2 \uparrow (k+9)$ as in Theorem 4.14. By Theorem 4.14, there is an i such that \mathcal{K}_i contains an arithmetic progression of length N . Now, the ‘‘jump’’ in the progression ℓ satisfies $\ell \leq P/N$. Therefore, $g(\ell, N) \leq \frac{1}{N} \leq \delta$. Since there are only finitely many \mathcal{K}_i 's, there exists a \mathcal{K}_i for which the $g(\ell, N)$ arithmetic progression condition is satisfied. \square

We end this section by recalling the following theorem of Bourgain and Tzafriri, which remains the strongest positive indicator that the answer to question 4.10 is yes. Recall that, for a set $K \subset \mathbb{Z}$, the density of K is defined to be

$$\text{dens}(K) = \lim_{n \rightarrow \infty} \frac{\#(K \cap [-n, n])}{2n},$$

provided that the limit exists.

Theorem 4.16. [1] *Let $U \subset \Pi$ be a set of positive measure. There exists a set $K \subset \mathbb{Z}$ such that $\text{dens}(K) > 0$ and $\{e^{2\pi i k x} \chi_U : k \in K\}$ is a Riesz basic sequence.*

In this section, we have shown that there are techniques available from the interplay of number theory with harmonic analysis that can be used to address the Feichtinger conjecture in the case of frames of translates. It is natural to wonder whether more sophisticated tools in this area can be used to settle the Feichtinger conjecture for frames of translates. Of particular interest would be to find the optimal functions $g(\ell, N)$ such that Theorem 4.13 and Corollary 4.15 hold. Indeed, if $g(\ell, N)$ in Theorem 4.13 could be chosen to be smaller than $g(\ell, N)$ in Corollary 4.15, then the Feichtinger conjecture would be disproven. There is currently a bit of a gap in the available estimates.

REFERENCES

- [1] J. Bourgain and L. Tzafriri, Invertibility of “large” submatrices with applications to the geometry of Banach spaces and harmonic analysis. *Israel J. Math.* **57** (1987), no. 2, 137–224.
- [2] J. Bourgain and L. Tzafriri, On a problem of Kadison and Singer. *J. Reine Angew. Math.* **420** (1991), 1–43.
- [3] M. Bownik, Anisotropic Hardy spaces and wavelets, *Mem. Amer. Math. Soc.* **164** (2003), no. 781, 122pp.
- [4] M. Bownik and K.-P. Ho, Atomic and Molecular Decompositions of Anisotropic Triebel-Lizorkin Spaces, *Trans. Amer. Math. Soc.* (to appear).
- [5] M. Bownik and D. Speegle, The Feichtinger conjecture for wavelet frames, frames of translates and frames of exponentials, *Can. J. Math.* (to appear).
- [6] P. Casazza, O. Christensen, N. Kalton, Frames of translates. *Collect. Math.* **52** (2001), no. 1, 35–54.
- [7] P. Casazza, O. Christensen, A. Lindner and R. Vershynin, Frames and the Feichtinger conjecture, *Proc. Amer. Math. Soc.* (to appear).
- [8] P. Casazza and R. Vershynin, Kadison-Singer meets Bourgain-Tzafriri, preprint. <http://www.math.missouri.edu/~pete/pdf/kadison-singer.pdf>
- [9] O. Christensen and A. Lindner, Decomposition of Riesz frames and wavelets into a finite union of linearly independent sets, *Lin. Alg. Appl.* **355** no. 1 (2002), 147–159.
- [10] M. Frazier, B. Jawerth, Decomposition of Besov spaces, *Indiana U. Math. J.* **34** (1985), 777–799.
- [11] M. Frazier, B. Jawerth, A Discrete Transform and Decomposition of Distribution Spaces, *J. Funct. Anal.* **93**, (1989), 34–170.
- [12] M. Frazier, B. Jawerth, G. Weiss, Littlewood-Paley Theory and the Study of Function Spaces, CBMS Regional Conference Ser., #79, American Math. Society (1991).
- [13] W. T. Gowers, A new proof of Szemerédi’s theorem, *Geom. Funct. Anal.*, Vol. 11 (2001), 465–588.
- [14] K. Gröchenig, Localized frames are finite unions of Riesz sequences, *Adv. Comput. Math.* **18** (2003), 149–157.
- [15] C. S. Güntürk, Approximating a bandlimited function using very coarsely quantized data: improved error estimates in sigma-delta modulation. *J. Amer. Math. Soc.* **17** (2004), no. 1, 229–242
- [16] H. Halpern, V. Kaftal and G. Weiss, Matrix pavings and Laurent operators. *J. Operator Theory* **16** (1986), no. 2, 355–374.
- [17] S. Jaffard, A density criterion for frames of complex exponentials *Michigan Math. J.* **38** (1991), 339–348.
- [18] L. Kuipers, H. Niederreiter, Uniform distribution of sequences. Pure and Applied Mathematics. Wiley-Interscience, New York-London-Sydney, 1974. xiv+390 pp.

- [19] P.-G. Lemarié-Rieusset, Projecteurs invariants, matrices de dilatation, ondelettes et analyses multi-résolutions, *Rev. Mat. Iberoamericana* **10**, (1994), 283–347.
- [20] H. L. Montgomery, Ten lectures on the interface between analytic number theory and harmonic analysis. CBMS Regional Conference Series in Mathematics, 84. *American Mathematical Society, Providence, RI*, 1994. xiv+220 pp.
- [21] H. L. Montgomery and R. C. Vaughan, Hilbert’s Inequality. *J. London Math. Soc. (2)* **8** (1974), 73–82.
- [22] A. Ron and Z. Shen, Weyl-Heisenberg frames and Riesz bases in $L^2(\mathbb{R}^d)$. *Duke Math. J.* **89** (1997), no. 2, 237–282.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403–1222
E-mail address: `mownik@uoregon.edu`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, SAINT LOUIS UNIVERSITY, 221
N. GRAND BLVD., ST. LOUIS, MO 63103
E-mail address: `speegled@slu.edu`